



# White root rot of avocado caused by Rosellinia necatrix

Fact Sheet (January 2023)

Hartley, J & van den Berg, N, Avocado Research Programme (FABI), University of Pretoria

# Background

Rosellinia necatrix Berl. ex Prill. is the causal agent of white root rot on a wide range of plant species. Spain, Israel and Japan have experienced severe R. necatrix infestation of avocado, grapevine, pear and apple orchards [1-4].

R. necatrix was first identified in South Africa in 1974 on apple and pear trees, and grapevines, in the Western Cape [5]. However, the pathogen was not eradicated and WRR was identified in South African avocado orchards in 2016 [6]. R. necatrix is now present on avocado in the Limpopo, Mpumalanga and KZN provinces.

#### Symptoms

White root rot is difficult to diagnose since foliar and root symptoms are unspecific, therefore, the disease is often mistaken for Phytophthora root rot. Primary symptoms include decline of vigour, retarded growth, and sparse foliage (Fig. A). Secondary symptoms include wilting of the leaves, chlorosis, leaf drop and death of branches [2]. Death can occur within a few weeks after the appearance of the foliar symptoms. However, some hosts do not show any foliar symptoms until the plant suddenly dies, sometimes with fruit and leaves still attached to the tree. Another distinguishing symptom is the presence of white mycelial growth on the root surface (Fig. B), in the soil (Fig. C) and underneath/on top of the bark at the crown of the tree (Fig. D). R. necatrix will spread to adjacent trees, with trees in close proximity to each other dying consecutively [2, 4].

# Biology

R. necatrix is an ascomycete fungus that can live as a saprophyte in the soil, allowing occupation of infected areas for extended periods. Mycelial growth can occur at temperatures ranging from 5-30°C, and at pH values between 6-8 in soil and even 4-9 in vitro [7].

# **Known Hosts**

This pathogen is widely distributed over five continents and has been reported on 347 different host species to date. Economically important hosts include apple, peach, orange, pear, grape, coffee, olive and avocado [8].

# Control

Control options are limited due to the hardy resting structures of the pathogen and deep soil penetration. Numerous common fungicides have been ineffective in eradicating R. necatrix. However, Fluazinam (BRAVE (fluazinam 500 g/l SC)) has shown potential in controlling white root rot from in vitro and in planta trials. An integrated approach of orchard sanitation, soil solarization, biocontrol (Trichoderma spp.) and chemical control has shown promising results in other countries [3, 9, 10]. Orchard sanitation includes immediate removal and burning of infected trees and restricting movement in the orchard. Solarization and good soil preparation prior to planting can also improve control. Avocado rootstocks in South Africa appear to be susceptible to white root rot.



Pictures provided by South African avocado growers and Jesse Hartley.

### What to do?

- 1. Monitor your trees for sudden, rapid decline and death and/or the presence of white mycelia on the trunk or in the soil.
- 2. Fill out a FABI diagnostic clinic form, available at app.informationhub.io/form/cl3cy217x00674ts66dcfz1nl or send an email to diagnostic.clinic@fabi.up.ac.za.
- Collect samples from the trunk (bark) and roots, package 3. separately in brown paper bags with a wet paper towel in each, place in a crate/polystyrene box and send to the FABI diagnostic clinic.

#### Contact Address

Avocado Research Programme, Prof. Noëlani van den Berg, Diagnostic clinic, FABI, 74 Lunnon street, University of Pretoria, Pretoria, 0083 Dr. Lieschen De Vos, +27 12 420 3938/5826.

### References

- 1. López-Herrera et al. 1998. Plant Disease 82: 1088-1092.
- 2. Pliego et al. 2012. Molecular Plant Pathology 13: 226-239.
- 3. Sztejnberg et al. 1987. Plant Disease 71: 365.
- 4. ten Hoopen & Krauss. 2006. Crop Protection 25: 89-107.
- 5. Van der Merwe & Matthee. 1974. Phytophylactica 6: 119-120.
- 6. van den Berg et al. 2018. Plant Disease 102: 1850.
- 7. Anselmi & Giorcelli. 1990. Forest Pathology 20: 175-183.
- 8. Farr et al. 2020. http://nt.ars-grin.gov/fungaldatabases/
- 9. Arjona-López & López-Herrera. 2020. BioControl 65: 1-9.
- 10. Arjona-López et al. 2019. Biological Control 136: 103996.