

# ***Sirex* species (Hymenoptera: Siricidae) and their parasitoids in *Pinus sylvestris* in eastern North America**

**Stefan J. Long**

Department of Entomology, Cornell University, Ithaca, New York 14853–2601, United States of America

**David W. Williams**

United States Department of Agriculture, Animal and Plant Health Inspection Service, Otis Laboratory, Building 1398, Otis Air National Guard Base, Massachusetts 02542, United States of America

**Ann E. Hajek<sup>1</sup>**

Department of Entomology, Cornell University, Ithaca, New York 14853–2601, United States of America

**Abstract**—Siricids and their parasitoids were reared from Scots pine (*Pinus sylvestris* L. (Pinaceae)) trees infested by *Sirex noctilio* F. in central New York State. *Sirex noctilio* accounted for 94.3% of the siricid specimens emerging, totaling 1313 specimens from six trees, with a maximum of 495 from one tree. Of the individuals emerging per tree,  $20.6 \pm 5.2\%$  were female. Two native siricids, *Sirex nigricornis* F. and *S. edwardsii* Brullé, also emerged from trees but in low numbers. Three hymenopteran parasitoid species that attack siricids emerged, totaling  $21.8 \pm 6.4\%$  parasitism per tree. *Ibalia leucospoides ensiger* Norton (Ibaliidae) was by far the most abundant parasitoid, at  $20.5 \pm 6.3\%$  parasitism per tree. The percentage of female *S. noctilio* emerging was positively correlated with wood diameter, whereas percent parasitism by *I. l. ensiger* was negatively correlated with wood diameter.

**Résumé**—Des siricides et leurs parasitoïdes ont été élevés à partir de pin sylvestre, *Pinus sylvestris* L. (Pinaceae), infestés par *Sirex noctilio* F. collectés dans le centre de l'état de New York. *Sirex noctilio* représentait 94,3 % des siricides émergés de six arbres, soit un total de 1313 spécimens, avec un maximum de 495 *S. noctilio* pour un seul arbre. Parmi les *S. noctilio* émergés,  $20,6 \pm 5,2\%$  étaient des femelles. Deux siricides natifs (*Sirex nigricornis* F. et *S. edwardsii* Brullé) ont également émergé des arbres, mais toujours en petit nombre. Trois hyménoptères parasitoïdes s'attaquant aux siricides émergèrent également, pour un taux de parasitisme global de  $21,8 \pm 6,4\%$  par arbre. *Ibalia leucospoides ensiger* Norton (Ibaliidae) était de loin le parasitoïde le plus abondant avec un taux de parasitisme moyen par arbre de  $20,5 \pm 6,3\%$ . Le pourcentage de femelles *S. noctilio* émergeant était positivement corrélé avec le diamètre du bois, tandis que le taux de parasitisme par *I. l. ensiger* était négativement corrélé avec le diamètre du bois.

The wood wasp *Sirex noctilio* F. (Hymenoptera: Siricidae) is native to Eurasia (Spradbery and Kirk 1978) but has been introduced accidentally to numerous countries in the Southern Hemisphere (Hurley *et al.* 2007). In North America, *S. noctilio* was first collected in New York State in 2004 (Hoebeker *et al.* 2005) and Ontario, Canada, in 2005 (de Groot 2007). In the United States of America, *S. noctilio* had been

reported from 29 counties in New York State, 7 counties in Pennsylvania, 4 counties in Michigan, and 1 county in Vermont by 2008 (National Agricultural Pest Information System (NAPIS) 2008).

In Southern Hemisphere countries where *S. noctilio* has been introduced, it has become a pest in commercial pine plantations by infesting and eventually killing pines (*Pinus* L. (Pinaceae)).

Received 26 September 2008. Accepted 5 February 2009.

<sup>1</sup>Corresponding author (e-mail: aeh@cornell.edu).  
doi: 10.4039/n08-068

Adult females inject a phytotoxic mucus and a symbiotic white rot fungus, *Amylostereum areolatum* (Fr.) Boidin, into trees when laying eggs; the mucus and fungus together kill the trees (Talbot 1977). Pines and *Sirex* spp. are not native to the Southern Hemisphere, and classical biological control programs have been widely used to introduce parasitic nematodes and hymenopterous parasitoids for control in *S. noctilio* (Hurley *et al.* 2007). These natural enemies were always collected from areas in the Palearctic and Nearctic regions where *S. noctilio* or other *Sirex* species are native.

*Sirex noctilio*, native North American species of *Sirex* L., and hymenopterous parasitoids (Ichneumonidae and Ibalidae) were studied in central New York State using naturally infested pine trees harvested in 2007. We investigated the species of hymenopterous parasitoids associated with *Sirex* species and explored the relationship between wood diameter and the abundance of *S. noctilio* and the most commonly occurring parasitoid, *Ibalia leucospoides ensiger* Norton (Ibalidae).

Six naturally infested Scots pine, *Pinus sylvestris* L., trees were sampled in late June 2007 at two sites in central New York State known to host populations of *S. noctilio*; in 2006 many trees at both sites exhibited symptoms of heavy attack (*i.e.*, resin beads) and at one site an average of >40 *S. noctilio* emerged from 2.1 m long sections of 30 infested trees (D.W. Williams, unpublished data). Four trees were felled in New Haven (43.47868°N, 76.35241°W) and two in Pompey (42.96212°N, 75.95803°W). For five trees, at least the lowest 1 m of each tree was not used, and for the additional tree, the lowest sample began 0.35 m above the ground; sampling was not designed to evaluate vertical distribution. Few insects emerged from one tree felled in Pompey from which wood samples with a total length of 150.5 cm and an average diameter of  $13.2 \pm 0.1$  cm (mean  $\pm$  SE) were obtained; these samples were not included in analyses of the association of emergence with wood diameter. The minimum diameter of wood used from the other trees was  $6.1 \pm 1.2$  cm and the maximum diameter  $17.3 \pm 3.3$  cm; the difference from smallest to largest diameter wood from a single tree was  $11.2 \pm 2.2$  cm. The total length of wood from each of these trees was  $10.0 \pm 1.1$  m. Wood was cut into bolts  $50.2 \pm 0.4$  cm (mean  $\pm$  SE) long and held at ambient temperature (18–30 °C) in cardboard barrels (69 cm height  $\times$  48 cm diameter) with screen lids. A contiguous section of a tree

was placed in each barrel for rearing insects, with  $5.2 \pm 0.5$  bolts/barrel (volume of wood/barrel =  $2.95 \pm 0.70$  m<sup>3</sup>), leaving space between bolts to allow insect emergence. From 2 July to 27 September 2007, all emerging *Sirex* spp. and parasitoids were collected every 1–3 days for identification. No more insects emerged and, on 27 October, the study was discontinued. The wood was not dissected to detect any remaining living or dead insects, so the total number of insects in the wood samples could have been underestimated. Voucher specimens of each *Sirex* and parasitoid species have been deposited in the Cornell University Insect Collection (Lot Number 1264).

The association between the average diameter of wood in a barrel and total number of adult *S. noctilio*/m<sup>3</sup> (both male and female) was analyzed by logistic regression using a Poisson distribution (Proc Genmod; SAS Institute Inc. 2007). To analyze the association between wood diameter and percent female *S. noctilio* or percent parasitism by *I. l. ensiger*, random intercept mixed models with trees as a random effect were used with arcsine square root transformed proportion of female *S. noctilio* or proportion of parasitism by *I. l. ensiger* (Proc Mixed; SAS Institute Inc. 2007).

*Sirex noctilio* emerged from wood in every barrel and accounted for 94.3% of the siricid specimens collected, totaling 1313 specimens from all samples (Table 1). Nearly two-thirds (66.1%) of *S. noctilio* emerged from wood with a diameter  $\leq 12.3$  cm. Of these specimens,  $20.6 \pm 5.2\%$  were female. Two native siricid species, *S. nigricornis* F. and *S. edwardsii* Brullé, also emerged from the same sections of the same trees as *S. noctilio*. This was somewhat surprising because both of these natives are known to carry the symbiotic fungus *Amylostereum chailletii* (Fr.) Boidin, whereas *S. noctilio* carries *A. areolatum* (Bedding and Akhurst 1978), and these fungal species are vegetatively incompatible (Thomsen and Koch 1999). Almost all *S. noctilio* (96.0%) emerged 8 days before either *S. nigricornis* or *S. edwardsii*. Most of the native *Sirex* (98.4% and 88.9% for *S. nigricornis* and *S. edwardsii*, respectively) emerged from wood with diameters ranging from 13.5 to 18.0 cm.

Three species of parasitoids emerged from *Sirex*-infested trees: two ichneumonids and one ibaliid (Table 2). The ibaliid (*I. l. ensiger*) accounted for 96.1% of all parasitoid specimens ( $91.7 \pm 45.1$  *I. l. ensiger* individuals/tree (mean  $\pm$

**Table 1.** Three *Sirex* species that emerged from six Scots pine (*Pinus sylvestris*) trees in central New York State in 2007.

	No./tree	Percentage of females/tree*	No. of trees infested
<i>S. noctilio</i>	218.8 (64.9)	20.6 (5.2)	6
<i>S. nigricornis</i>	10.2 (8.0)	58.9 (16.1)	2
<i>S. edwardsii</i>	3.0 (2.4)	6.7 (6.7)	3

**Note:** Values are shown as the mean with SE in parentheses.

\*Means were calculated using trees in which the species was present.

**Table 2.** Three parasitoid species associated with *Sirex* spp. emerging from six *Pinus sylvestris* trees in central New York State in 2007.

	No./tree	Percent parasitism/tree*	Percentage of females/tree†	No. of trees present
<i>Ibalia leucospoides ensiger</i>	91.7 (45.1)	20.5 (6.3)	45.7 (9.7)	5
<i>Rhyssa lineolata</i>	2.5 (1.2)	0.8 (0.5)	62.5 (37.5)	4
<i>Megarhyssa nortoni</i>	1.2 (0.7)	0.5 (0.3)	62.4 (9.0)	3

**Note:** Values are shown as the mean with SE in parentheses.

\*Calculated as combined parasitism of *S. noctilio*, *S. nigricornis*, and *S. edwardsii*.

†Means were calculated using trees in which the species was present.


SE)) emerging from five of the six trees. The two ichneumonids, *Rhyssa lineolata* (Kirby) and *Megarhyssa nortoni* (Cresson), emerged from only four and three trees, respectively, with a maximum of seven and three individuals per tree, respectively. Both ichneumonid species emerged from both sites. In one section of one tree, both *I. l. ensiger* and *R. lineolata* emerged along with *S. noctilio* and both native siricids. *Megarhyssa nortoni* did not emerge from the same tree sections as the native siricids, though *M. nortoni* emerged from one of the trees from which *S. edwardsii* emerged. Because all three *Sirex* species emerged from the trees that were sampled, it is not possible to calculate percent parasitism for each *Sirex* species separately. Parasitism of *Sirex* spp. by all parasitoids was  $21.8 \pm 6.4\%$  per tree.

No relationship was found between the total density of *S. noctilio* and the average diameter of wood. However, a significantly larger percentage of female *S. noctilio* emerged with increasing wood diameter (arcsine square root (proportion of females) =  $-0.04408 + 0.04748 \times$  wood diameter;  $F_{1,16.7} = 6.59$ ,  $P = 0.0202$ ) (Fig. 1). The inverse relationship was found for percent parasitism by *I. l. ensiger*, with fewer parasitoids emerging as wood diameter increased (arcsine square root (proportion of parasitism) =  $0.9975 - 0.03981 \times$  wood diameter;  $F_{1,16} = 8.01$ ,  $P = 0.0121$ ) (Fig. 1).

The parasitoids reared in this study could have parasitized any of the three siricid species reared, but we hypothesize that much of the parasitism likely occurred on *S. noctilio*. Our study sites were 27.5 and 54.0 km from the site where *S. noctilio* was first detected in 2004, and *S. noctilio* has been found killing pines at these study sites since 2006 (D.W. Williams, unpublished data). Although the parasitoids reared in this study are Nearctic natives and *S. noctilio* is native to the Palearctic Region, parasitoids associated with concealed hosts frequently have broad host ranges (Strand 1986). In fact, two of the three parasitoid species that emerged in this study have previously been used for classical biological control of *S. noctilio* in the Southern Hemisphere (Hurley *et al.* 2007), and all three species have been recorded parasitizing numerous siricid species (Krombein *et al.* 1979). In our study, *I. l. ensiger*, which parasitizes eggs and younger larvae (Murphy 1998), was much more abundant than the two rhyssine ichneumonids (*M. nortoni* and *R. lineolata*), which attack later instars (Murphy 1998). In contrast, in Tasmania, parasitism of *S. noctilio* by rhyssines was much more common than parasitism by *I. leucospoides* (Taylor 1978).

Chiarello Penteado *et al.* (2000) showed that most *S. noctilio* and most *I. leucospoides* in Brazil emerged from the central 50% of trees, based on tree diameter. Our results showed that

**Fig. 1.** Percentage of female *Sirex noctilio* (solid line) and percent parasitism by *Ibalia leucospoides ensiger* (dashed line) emerging from tree sections of various diameters; lines were calculated using arcsine square root transformed proportions. Only 10 male *S. noctilio* and no parasitoids emerged from wood samples totaling 150.5 cm in length from one tree in Pompey, New York; samples from this tree were not included in statistical analyses.



although the total number of *S. noctilio* emerging did not vary with tree diameter, the percentage of females increased significantly with tree diameter, which is similar to observations of Morgan and Stewart (1966) and Hurley *et al.* (2008). In contrast, the percentage of hosts parasitized by *I. l. ensiger* decreased significantly as tree diameter increased. This allocation of a greater percentage of female *S. noctilio* to larger diameter sections of the tree could allow female offspring to escape parasitism by *I. l. ensiger*. We hypothesize that oviposition by *I. l. ensiger* could be limited by some feature of larger diameter wood (*e.g.*, bark thickness) or by height in the tree, but these factors require further investigation.

We thank K. Zylstra, C. Tait, and M. Crawford for their help in the field. E.R. Hoebeke of the Cornell University Insect Collection provided species identifications. B. Hurley and E.A. Cameron both assisted with information about sources of *S. noctilio* parasitoids introduced to the Southern Hemisphere. F. Vermeylen assisted with statistical

analyses. G. Desurmont provided the French translation of the abstract. Reviews by J.K. Lieberr and an anonymous reviewer substantially improved the manuscript. This study was funded through a cooperative agreement with the United States Department of Agriculture, Animal and Plant Health Inspection Service.

Bedding, R.A., and Akhurst, R.J. 1978. Geographical distribution and host preferences of *Deladenus* species (Nematoda: Neotylenchidae) parasitic in siricid woodwasps and associated hymenopterous parasitoids. *Nematologica*, **24**: 286–294.

Chiarello Penteado, S.R., Batista de Oliveira, E., and Iede, E.T. 2000. Distribuição da vespa-da-madeira e de seus inimigos naturais ao longo do tronco de *Pinus*. *Boletim de Pesquisa Florestal*, **40**: 23–34.

de Groot, P. 2007. An overview of the *Sirex noctilio* situation in Canada [online]. In *Proceedings of the International Sirex Symposium*, 10 and 11 May 2007, Pretoria, South Africa [online]. p. 17. Available from <http://src.fabinet.up.ac.za/sirex/pdf/SirexsymposiumproceedingsFULLfin.pdf> [accessed 16 July 2008].

- Hoebeke, E.R., Haugen, D.A., and Haack, R. 2005. *Sirex noctilio*: discovery of a Palearctic siricid wasp in New York. Newsletter of the Michigan Entomological Society, **50**(1, 2): 24–25.
- Hurley, B.P., Slippers, B., and Wingfield, M.J. 2007. A comparison of control results for the alien invasive woodwasp, *Sirex noctilio*, in the southern hemisphere. Agricultural and Forest Entomology, **9**: 159–171.
- Hurley, B.P., Slippers, B., Croft, P.K., Hatting, H.J., van der Linde, M., Morris, A.R., Cyer, C., and Wingfield, M.J. 2008. Factors influencing parasitism of *Sirex noctilio* (Hymenoptera: Siricidae) by the nematode *Deladenus siricidicola* (Nematoda: Neotylenchidae) in summer rainfall areas of South Africa. Biological Control, **45**: 450–459.
- Krombein, K.V., Hurd, P.D., and Smith, D.R. 1979. Catalog of Hymenoptera in America North of Mexico. Smithsonian Institution Press, Washington, D.C.
- Morgan, F.D., and Stewart, N.C. 1966. The biology and behavior of the woodwasp *Sirex noctilio* F. in New Zealand. Transactions of the Royal Society of New Zealand, **7**: 195–204.
- Murphy, S.T. 1998. Indigenous siricid spp. parasitoid communities and principal biological control agents of *Sirex noctilio* in Australasia: a review. United States Department of Agriculture Forest Service FHTET 98-13. pp. 31–35.
- National Agricultural Pest Information System, Center for Environmental and Regulatory Information Systems. 2008. Reported status of European woodwasp — *Sirex noctilio* [online]. Available at <http://pest.ceris.purdue.edu/searchmap.php?selectName=ISBBADA> [accessed 15 March 2009].
- SAS Institute Inc. 2007. SAS® for Windows. Version 9.1.3. SAS Institute Inc., Cary, North Carolina.
- Spradbery, J.P., and Kirk A.A. 1978. Aspects of the ecology of siricid woodwasps (Hymenoptera: Siricidae) in Europe, North Africa and Turkey with special reference to the biological control of *Sirex noctilio* F. in Australia. Bulletin of Entomological Research, **68**: 341–359.
- Strand, M.R. 1986. The physiological interactions of parasitoids with their hosts and their influence on reproductive strategies. In Insect parasitoids. Edited by J. Waage and D. Greathead. Academic Press, London. pp. 97–136.
- Talbot, P.H.B. 1977. The *Sirex*–*Amylostereum*–*Pinus* association. Annual Review of Phytopathology, **15**: 41–54.
- Taylor, K.L. 1978. Evaluation of the insect parasitoids of *Sirex noctilio* (Hymenoptera: Siricidae) in Tasmania. Oecologia, **32**: 1–10.
- Thomsen, I.M., and Koch, J. 1999. Somatic compatibility in *Amylostereum areolatum* and *A. chailletii* as a consequence of symbiosis with siricid woodwasps. Mycological Research, **103**: 817–823.