

Host Range of *Fusarium* Dieback and Its Ambrosia Beetle (Coleoptera: Scolytinae) Vector in Southern California

Akif Eskalen, Department of Plant Pathology and Microbiology, Richard Stouthamer, Department of Entomology, Shannon Colleen Lynch, Department of Plant Pathology and Microbiology, and Center for Conservation Biology, Paul F. Rugman-Jones, Department of Entomology, and Mathias Twizeyimana and Alex Gonzalez, Department of Plant Pathology and Microbiology, University of California, Riverside, 92521; and Tim Thibault, Huntington Library, Art Collections, and Botanical Gardens, San Marino 91108

Abstract

Eskalen, A., Stouthamer, R., Lynch, S. C., Twizeyimana, M., Gonzalez, A., and Thibault, T. 2013. Host range of *Fusarium* dieback and its ambrosia beetle (Coleoptera: Scolytinae) vector in southern California. *Plant Dis.* 97:938-951.

The polyphagous shot hole borer (PSHB) is an invasive ambrosia beetle that forms a symbiosis with a new, as-yet-undescribed *Fusarium* sp., together causing *Fusarium* dieback on avocado and other host plants in California and Israel. In California, PSHB was first reported on black locust in 2003 but there were no records of fungal damage until 2012, when a *Fusarium* sp. was recovered from the tissues of several backyard avocado trees infested with PSHB in Los Angeles County. The aim of this study was to determine the plant host range of the beetle–fungus complex in two heavily infested botanical gardens in Los Angeles County. Of the 335 tree species observed, 207 (62%), representing 58 plant families, showed signs and symptoms consistent

with attack by PSHB. The *Fusarium* sp. was recovered from 54% of the plant species attacked by PSHB, indicated by the presence of the *Fusarium* sp. at least at the site of the entry hole. Trees attacked by PSHB included 11 species of California natives, 13 agriculturally important species, and many common street trees. Survey results also revealed 19 tree species that function as reproductive hosts for PSHB. Additionally, approximately a quarter of all tree individuals planted along the streets of southern California belong to a species classified as a reproductive host. These data suggest the beetle–disease complex potentially may establish in a variety of plant communities locally and worldwide.

Since the beginning of 2012, an emerging plant disease, referred to as *Fusarium* dieback (FD), has been detected on several host plants in Los Angeles and Orange Counties, CA (7). FD is caused by the combined effects of a tiny (<2.5 mm) invasive ambrosia beetle (Scolytinae) and its symbiotic fungal partner, a currently undescribed *Fusarium* sp. (7,15). Adult female beetles burrow into a tree, creating brood galleries beyond the cambium, which are concurrently inoculated with the *Fusarium* sp. The fungus colonizes the gallery walls and becomes the sole source of food for developing larvae and adult beetles (7,15). The fungus invades the vascular tissue of the tree, blocking water and nutrients from the roots to the rest of the tree, eventually causing branch dieback and tree death (Fig. 1) (7,15).

The specific identity of the beetle is currently unknown. It is morphologically indistinguishable from the tea shot hole borer (TSHB), *Euwallacea fornicatus*, a serious pest of tea (*Camellia sinensis*) in Sri Lanka and India (5) (Fig. 2A–C). However, there are significant differences in nuclear and mitochondrial DNA sequences between the California beetles and those from tea plantations in Sri Lanka and other Asian collection sites (P. F. Rugman-Jones and R. Stouthamer, unpublished data). This indicates that the California beetle is most likely a separate, congeneric species (representative mitochondrial sequences for the two species are available in GenBank [1], accession numbers JX912723–JX912725). Species within the genus *Euwallacea* have been subject to much revision and synonymization (28); therefore, establishing the exact identity of this beetle is likely to require extensive taxonomic research. Thus, for the purpose of this study, the beetle is simply

referred to as *Euwallacea* sp. and the common name polyphagous shot hole borer (PSHB) is used.

In California, PSHB was first collected in 2003 from several tree species (20) but there were no reports of fungal damage before 2012. Preliminary survey data in 2012 showed that backyard avocado trees and many other woody plants in the urban landscape, especially castor bean (*Ricinus communis*), were heavily infested with this beetle–fungus disease complex in Los Angeles and Orange counties (A. Eskalen, unpublished data). Some of the most heavily infested areas in southern California are home to two of its biggest botanical gardens, the Los Angeles (LA) Arboretum and The Huntington Library, Art Collections and Botanical Gardens, which are relatively close to the original 2003 find of the beetle. These botanical gardens harbor a wide range of plant species that represent unique and common ecosystems from all over the world, and contain all the host species planted throughout urban forests in

Corresponding author: A. Eskalen, E-mail: akif.eskalen@ucr.edu

*The e-Xtra logo stands for “electronic extra” and indicates that Figures 1 to 5, 7, and 8 appear in color in the online edition.

Accepted for publication 24 January 2013.

<http://dx.doi.org/10.1094/PDIS-11-12-1026-RE>
© 2013 The American Phytopathological Society




Fig. 1. Wood discoloration and xylem vessel plugging caused by *Fusarium* sp. on *Albizia julibrissin*.

southern California. In addition, the collections receive more consistent and even care as opposed to lands with multiple management strategies. Therefore, the botanical gardens provided a perfect natural laboratory to investigate the plant host range of the beetle–fungus complex. An understanding of the host range of the beetle–fungus complex is important in determining the present disease distribution and to make predictions about its future distribution.

At present, the beetle–fungus complex appears to be limited to the urban forest. In itself, this will likely present economic problems for households and cities that must bear the cost of removing dead or dying trees. However, the same disease complex was also found in Israel in 2009 in commercial avocado orchards, where it causes significant damage to avocado trees (15). Thus, the presence of FD in urban landscapes throughout southern California is a concern for the California avocado industry because it may eventually spread to commercial avocado groves. Similarly, the wide host range of this beetle may ultimately threaten native trees in the natural forests of southern California.

The objectives of this study were to (i) determine the plant host range of the PSHB and *Fusarium* sp. and (ii) characterize their symptoms on different host plants.

Materials and Methods

Host identification and symptom characterization. Trees were visited and identified to species at the LA Arboretum and The Huntington Library, Art Collections and Botanical Gardens in Los Angeles County; specific locations of sampled trees were noted using global positioning system equipment (Garmin 60CSx, 5- to 10-m accuracy). Each botanical garden covers approximately 48 ha and, in general, the species collections housed in each are different. Although the gardens contain a large number of plant species, the criteria in host plant selection included (i) common street trees found in southern California (4), (ii) agriculturally important trees and vines, (iii) species of tree families, and (iv) trees randomly encountered in the search for species outlined using the criteria i–iii.

Signs and symptoms of FD (presence or absence; see Results), severity of beetle colonization (0 = none, 1 = minor [1 to 10 entry holes], 2 = moderate [10 to 30 entry holes], 3 = [>31 entry holes]), and branch dieback (0 = none, 1 = minor [1 to 25% of branches showed dieback], 2 = moderate [25 to 50%], and 3 = severe [>50%]) were also recorded. For each tree species showing symptoms characteristic of FD (one tree per species), tissue samples were collected by extracting symptomatic tissue from the trunk with a sterilized knife to a depth beyond the cambium. Samples were returned to the laboratory (University of California, Riverside) for morphological and molecular identification of the beetle, the fungus, or both. If visible, beetles were collected and put directly into 95% ethanol for later molecular identification (see below). The data were analyzed using JMP 9.0.0 statistical software (SAS 2012). Logistic regression was used to determine the presence and absence of *Fusarium* sp., gumming, and severity of beetle attack as contributing factors to branch dieback >50% at $\alpha = 0.05$.

The suitability of a tree species to act as a reproductive host for the PSHB was evaluated, where possible, by cutting into trees and

confirming the presence of eggs, larvae, pupae, or teneral females or males in the galleries. Trees that did not show signs of heavy infestation by PSHB were not cut to examine and confirm as a reproductive host because species of these trees were limited in number within the botanical gardens' collections.

***Fusarium* sp. and PSHB confirmation.** To carry out fungal isolations, gross contamination was removed before bringing plant samples into a clean area by briefly flaming samples and removing the outer surface with a sterilized paring knife. All samples were then split in half and pieces excised from the leading margin of clean necrotic tissue were plated onto potato dextrose agar (Difco) amended with 100 μg of tetracycline (Fisher Scientific). Plates were incubated at 25°C, and fungal growth was subcultured for identification and long-term storage after 7 to 10 days.

Total genomic DNA was extracted from the mycelium of each isolate using the method described by Ceniz (2). Portions of the following four gene fragments were chosen for multilocus sequence typing based on published genetic analyses (3,16–18,29): the internal transcribed spacer (ITS) region, translation elongation factor 1- α (EF1- α), the second-largest subunit of RNA polymerase (*RPB2*), and DNA-directed RNA polymerase subunit 1 (*RPB1*). Each 25- μl polymerase chain reaction (PCR) mixture contained 19.4 μl of PCR-grade water, 2.5 μl of standard Taq reaction buffer, 0.5 μl of dNTPs (10 mM), 0.125 μl of NEB *Taq* DNA polymerase (5,000 U/ml; New England BioLabs), 0.5 μl of each primer at 10 mM, and 1.5 μl of template DNA. PCR was carried out in a thermal cycler (Bio-Rad Laboratories, Inc.) using published cycling conditions (14,17,25). Amplification products were separated by electrophoresis in 1.5% agarose gels in 1.0 \times Tris-boric acid-EDTA buffer and visualized under UV light after staining with SYBR Green (Invitrogen). PCR products were purified using an Exo SAP-IT kit (Affymetrix). ITS, EF1- α , RPB1, and RPB2 regions were sequenced in both directions at the Institute for Integrative Genome Biology at the University of California, Riverside. Sequences of isolates from the present study were confirmed as *Fusarium* sp. after a BLAST search comparison in GenBank, which showed they were 100% homologous to known sequences of isolates of *Fusarium* sp. from Eskalen et al. and Mendel et al. (7,15). The ITS, EF1- α , RPB1, and RPB2 sequences used in this study were deposited into GenBank.

PSHB collection and confirmation. When present, live beetles were collected directly into 95% ETOH and returned to the laboratory for identification. In the field, beetles were collected from the surface of the trees using forceps. If they were in a gallery, the beetles were first coaxed to the entrance by spraying 95% ethanol in the gallery and then removed with the forceps (11).

In the laboratory, DNA was extracted from individual beetles by grinding two to three tibia in 2 μl of proteinase-K (>600 mAU ml⁻¹; Qiagen) and 35 μl of Chelex 100 resin (Bio-Rad Laboratories) followed by incubation for 1 h at 55°C, then for 10 min at 99°C. A section of the cytochrome c oxidase subunit I gene commonly used in barcoding studies (10) was amplified using the primers LCO1490 (5'-GGTCAACAAATCATAAAGATATTGG-3') and



Fig. 2. Polyphagous shot hole borer (PSHB) (*Euwallacea* sp.): A, female; B, male; C, size comparison of both male (left) and female (right) on a penny.

Table 1. Hosts of the polyphagous shot hole borer and their susceptibility to Fusarium dieback (FD)^a

Family, species ^b	Common name	N ^c	Crop ^d	Origin	FD ^e	S	P	G
Aceraceae s.s. (Sapindaceae s.l.)								
* <i>Acer buergerianum</i>	Trident maple	4	...	East China, Taiwan	Y
<i>A. davidii</i>	Pere David's maple	China	...	Y
<i>A. caudatifolium</i> (syn. <i>A. kawakamii</i>)	Kawakami maple	Taiwan	...	Y
* <i>A. macrophyllum</i>	Big leaf maple	Western North America	Y	Y
* <i>A. negundo</i>	Box elder	North America	Y	Y
* <i>A. palmatum</i> 'Bonfire'	Bonfire Japanese maple	East Asia	Y
* <i>A. paxii</i>	Evergreen maple	3	...	Southwest China	Y	Y
<i>A. × freemanii</i>	Freeman maple	Eastern North America	...	Y
Alangiaceae s.s. (Cornaceae s.l.)								
<i>Alangium chinensis</i>	Tropical Africa to China	Y	Y
Anacardiaceae								
<i>Pistacia chinensis</i>	Chinese pistache	74	...	China	Y
<i>Schinus terebinthifolius</i>	Brazilian pepper tree	333	...	Brazil	Y	Y
Apocynaceae								
<i>Thevetia thevetioides</i>	Yellow oleander	Mexico	...	Y
Aquifoliaceae								
<i>Ilex aquifolium</i>	English holly	Western and Southern Europe, Northwest Africa, Southwest Asia
* <i>I. cornuta</i>	Chinese holly	China and Korea	Y	Y	...	Y
<i>I. latifolia</i>	Tarajo holly	Japan and China	Y	Y	Y	...
Araliaceae								
<i>Cussonia spicata</i>	Spiked cabbage tree	South Africa	Y	Y	...	Y
<i>Fatsia japonica</i>	Japanese aralia	Japan	Y	Y	...	Y
Arecaceae								
<i>Brahea armata</i>	Mexican blue palm	1	...	Western Mexico
<i>Butia capitata</i>	Pindo palm	Brazil, Uruguay, Argentina	...	Y	...	Y
<i>Livistona chinensis</i>	Chinese fan palm	Australia, Southeast Asia	Y	Y
<i>Washingtonia filifera</i>	Desert fan palm	54	...	Southwest United States, Northwest Mexico	Y	Y
Asparagaceae								
<i>Dracaena draco</i>	Canary islands dragon tree	Canary Islands, Cape Verde, Madeira, Morocco	...	Y
Asteraceae								
<i>Verbesina gigantea</i>	Crownbeard	Tropical America	Y
Betulaceae								
<i>Alnus incana</i>	Grey alder	Europe, Caucasus	...	Y	...	Y
<i>A. rhombifolia</i>	white alder	72	...	Western United States	...	Y
<i>Betula pendula</i>	Silver birch	12	...	Europe, Asia Minor	Y
<i>Corylus colurna</i>	Turkish hazelnut	...	Y	Southeast Europe, Asia Minor, Western Asia	Y
Bignoniaceae								
<i>Catalpa speciosa</i>	Northern catalpa	8	...	Central and Eastern United States	Y	Y
<i>Handroanthus impetiginosus</i> (syn. <i>Tabebuia impetiginosa</i>)	Brazilwood	Mexico to South America	Y	...
Bombacaceae s.s. (Malvaceae s.l.)								
<i>Bombax ceiba</i>	Cotton tree	India, China, Southeast Asia	Y
<i>Ceiba speciosa</i>	Silk floss tree	14	...	Brazil, Argentina	Y	Y	...	Y
<i>Pseudobombax ellipticum</i>	Shaving brush tree	Mexico and Central America	Y
Buddlejaceae								
<i>Nuxia floribunda</i>	Forest elder	Africa
Bursaceae								
<i>Bursera hindsiana</i>	Western Mexico	...	Y
Casuarinaceae								
<i>Casuarina cunninghamiana</i>	She-oak	Northeast Australia	...	Y
Cunoniaceae								
<i>Cunonia capensis</i>	Buttetspoon tree	South Africa	...	Y
Cornaceae								
<i>Cornus controversa</i>	Giant dogwood	Himalayas to Japan	Y
Cupressaceae								
<i>Juniperus chinensis</i>	Chinese juniper	26	...	Northeast Asia	Y	Y
<i>J. virginiana</i>	Eastern red-cedar	Eastern North America	Y
Ebenaceae								
<i>Diospyros kaki</i>	Japanese persimmon	2	Y	Asia	...	Y
<i>D. lycidioides</i>	Y	South Africa	Y	Y
Elaeocarpaceae								
<i>Crinodendron patagua</i>	Lily of the valley tree	Chile	...	Y
Ericaceae								
<i>Arbutus unedo</i>	Strawberry tree	Southwest Ireland to Asia Minor	...	Y

(continued on next page)

^a S = staining, P = powder, G = gumming, and Y = "yes".^b Asterisks: * indicates reproductive hosts and ** indicates critically endangered tree.^c Number of times planted along streets of City of Orange (4).^d Agriculturally important crop.^e FD susceptible.

Table 1. (continued from preceding page)

Family, species ^b	Common name	N ^c	Crop ^d	Origin	FD ^e	S	P	G
Euphorbiaceae								
<i>Aleurites fordii</i>	Tung oil tree	Asia	Y
<i>Bischofia javanica</i>	Bishop wood	Tropical Asia	Y	Y
<i>Jatropha</i> cf. <i>cinerea</i>	Limberbush	Arizona, Western Mexico	Y
<i>Manihot esculenta</i>	Cassava	Brazil
* <i>Ricinus communis</i>	Castor bean	Mediterranean basin, East Africa, India	Y	Y
<i>Sapium sebiferum</i>	Chinese tallow tree	China, Japan	Y	Y
Fabaceae								
<i>Acacia caven</i>	Roman cassie	South America	...	Y
<i>A. stenophylla</i>	Shoestring acacia	Australia	Y
<i>A. visco</i>	Arca	Argentina	Y
<i>Albizia kalkora</i>	Kalkora mimosa	Southern Asia	...	Y
* <i>A. julibrissin</i>	Silk tree	Southwest to east Asia	Y	Y
<i>Bauhinia</i> × <i>blakeana</i>	Orchid tree	1	...	China	Y	Y
<i>B. galpinii</i>	Red orchid bush	South Africa	...	Y	Y	...
<i>Calpurnia aurea</i>	Southern India, Africa	Y	Y	...	Y
<i>Cassia brewsteri</i>	Cassia pea	Northeast Australia	Y	Y	...	Y
* <i>Castanospermum australe</i>	Black bean	Northeast Australia, New Caledonia	Y	Y	...	Y
* <i>Cercidium floridum</i>	Blue palo verde	Southwest United States, Western Mexico	Y	Y
* <i>C. × sonorae</i>	Brea	Northwest Mexico	Y	Y
<i>Cladrastris sinensis</i>	Chinese yellow wood	China	...	Y	...	Y
<i>Erythrina folkersii</i>	Coral tree	Central America	Y	...	Y	...
<i>E. coralloides</i>	Flame coral tree	Arizona and Mexico	...	Y
* <i>Erythrina corallodendron</i>	Coral tree	West Indies	Y	Y
<i>E. crista-galli</i>	Ceibo	South America	Y	Y	...	Y
<i>E. humeana</i>	Dwarf coral tree	South Africa, Mozambique	Y	Y	Y	...
<i>E. lysistemon</i>	Common coral tree	South Africa	Y	Y	Y	...
<i>E. × sykesii</i>	Coral tree	Horticultural origin	Y	Y
<i>Inga feuilleii</i>	Pacay	South America	Y	Y
<i>Lysiphyllum carronii</i>	Queensland ebony	Australia	Y	Y
* <i>Parkinsonia aculeata</i>	Palo verde	Tropical America	Y	Y
<i>Pithecellobium</i> sp.	Y
<i>Schotia brachypetala</i>	Huilboerboon	Southern Africa	Y	Y	...	Y
<i>Senna racemosa</i>	Limestone senna	Central and South America	...	Y
<i>Senna racemosa</i> var. <i>liebmanni</i>	Central and South America	Y	Y
<i>S. spectabilis</i> var. <i>spectabilis</i>	Y	Y	Y
<i>Styphnolobium japonicum</i> (syn. <i>Sophora japonica</i>)	Pagoda tree	East Asia	Y
<i>Tipuana tipu</i>	Tipu tree	1	...	Brazil, Argentina, Bolivia	...	Y
<i>Wisteria sinensis</i>	Chinese wisteria	China	Y
<i>W. floribunda</i> 'Macrobotyrus'	Japanese wisteria	Japan	Y	Y
<i>Zenia insignis</i>	South China and Vietnam	...	Y
Fagaceae								
<i>Fagus sylvatica</i>	European beech	Central Europe to Caucasus	Y	Y
<i>F. sylvatica</i> 'Riversii'	Riversii European Beech	Horticultural origin	Y	...
<i>Quercus acutissima</i>	Sawtooth oak	China, Korea and Japan
* <i>Q. agrifolia</i>	Coast live oak	13	...	California, Baja California	Y	Y
<i>Q. alba</i>	White oak	Eastern North America
<i>Q. chrysolepis</i>	Canyon live oak	Western North America	Y	Y
<i>Q. engelmannii</i>	Engelmann Oak	Southwest California	...	Y
<i>Q. ilex</i>	Holly oak	306	...	Mediterranean	Y	Y
<i>Q. lobata</i>	Valley oak	California	Y	Y
<i>Q. macrocarpa</i>	Bur oak	Central and Eastern North America	Y	Y
<i>Q. mexicana</i>	Coahuatl	Mexico	Y	Y	...	Y
* <i>Q. robur</i>	English oak	Europe, Anatolia, and North Africa	Y	Y
<i>Q. rubra</i>	Northern red oak	7	...	Eastern North America	...	Y	Y	Y
<i>Q. suber</i>	Cork oak	Southwest Europe and Northwest Africa	Y	Y
<i>Q. virginiana</i>	Southern live oak	Southeastern United States	Y	Y
Hamamelidaceae s.l. (Altingiaceae s.s.)								
<i>Liquidambar formosana</i>	Chinese sweet gum	12	...	South China, Taiwan	Y	Y
* <i>L. styraciflua</i>	American sweetgum	4,916	...	Eastern North America	Y	Y
Hydrophyllaceae s.s. (Boraginaceae s.l.)								
<i>Wigandia urens</i>	Peru	...	Y
Juglandaceae								
<i>Carya illinoensis</i>	Hickory	36	...	South-Central North America	Y	Y	...	Y
<i>Juglans mandshurica</i> (syn. <i>Juglans formosana</i>)	Hayata	China, Taiwan	Y	...
<i>J. nigra</i>	Eastern black walnut	5	Y	Eastern North America	...	Y
<i>Pterocarya</i> sp.	Wingnut tree	Asia	...	Y	...	Y

(continued on next page)

Table 1. (continued from preceding page)

Family, species ^b	Common name	N ^c	Crop ^d	Origin	FD ^e	S	P	G
Lauraceae								
<i>Cinnamomum glanduliferum</i>	False camphor tree	China
<i>C. camphora</i>	Camphor tree	690	...	Japan to Tropical Asia	Y	Y	Y	...
<i>Nothaphoebe cavaleri</i>	South China	...	Y
* <i>Persea americana</i>	Avocado	70	Y	Mexico, Guatemala, West Indies	Y	Y	Y	...
<i>Machilus thunbergii</i> (syn. <i>Persea thunbergii</i>)	Asian avocado	East Asia	Y	Y
<i>Umbellularia californica</i>	California bay laurel	Western North America	Y	Y
Magnoliaceae								
<i>Magnolia</i> 'Columbus'	Columbus magnolia	Horticultural origin	...	Y
<i>M. delavayi</i>	Chinese evergreen magnolia	Southwest China	...	Y
<i>M. doltsopa</i>	Sweet michelia	Southwest China, Himalayas	...	Y
<i>M. grandiflora</i>	Southern magnolia	2,174	...	Southeast United States	Y	Y
<i>M. guatemalensis</i>	Mamey	Central America	...	Y	...	Y
<i>M. × soulangeana</i>	Saucer magnolia	8	...	Horticultural origin	...	Y	...	Y
<i>M. × veitchii</i>	Veitch's magnolia	Horticultural origin	Y	Y
Malvaceae								
<i>Dombeya cacuminum</i>	Strawberry tree	Madagascar	Y	...	Y	...
<i>Brachychiton acerifolius</i>	Illawarra flame tree	46	...	Northeast Australia	Y	Y	...	Y
<i>B. australis</i>	Kurrajong	Northern Australia	Y	Y
<i>B. discolor</i>	lacebark tree	Northeast Australia	Y
<i>B. rupestris</i>	Queensland bottle tree	Northeast Australia	Y	Y
<i>Chiranthodendron pentadactylon</i>	Devil's hand tree	Mexico and Central America	...	Y	...	Y
<i>Firmiana simplex</i>	Chinese parasol tree	East Asia	Y
Meliaceae								
<i>Melia azedarach</i>	Cape lilac	11	...	India	...	Y
<i>Swietenia chickrassa</i> (syn. <i>Chukrasia tabularis</i>)	India and China	Y
Melianthaceae								
<i>Melianthus major</i>	Giant honey flower	South Africa	Y
Menispermaceae								
<i>Cocculus laurifolius</i>	Laurel-leaved snail tree	Himalayas to Japan	Y	Y
<i>C. orbiculatus</i> (syn. <i>C. trilobus</i>)	Korean moonseed	East and Southeast Asia	Y	Y	...	Y
Monimiaceae								
<i>Peumus boldus</i>	Boldo	Chile	...	Y
Moraceae								
<i>Broussonetia papyrifera</i>	Paper mulberry	Eastern Asia	...	Y
<i>Ficus benjamina</i>	Weeping fig	1,182	...	Southeast Asia and Australia	...	Y
<i>F. macrophylla</i>	Moreton Bay fig	2	...	Northeast Australia	Y	Y	...	Y
<i>F. maxima</i>	Fig	Tropical America
<i>F. platypoda</i>	Desert rock fig	Australia and Indonesia	Y	Y	...	Y
<i>Morus alba</i>	White mulberry	43	Y	China	Y	Y
Myrtaceae								
<i>Callistemon salignus</i>	Willow bottlebrush	Australia
<i>C. viminalis</i>	Weeping bottlebrush	91	...	Australia	...	Y
<i>Eucalyptus camaldulensis</i>	River red gum	2	...	Australia
<i>E. cinerea</i>	Argyle apple	Southeast Australia	...	Y
<i>E. ficifolia</i>	Red flowering gum	33	...	Western Australia	...	Y	...	Y
<i>E. froggattii</i>	Southeast Australia	...	Y	...	Y
<i>E. kitsoniana</i>	Gippsland mallee	Southeast Australia	...	Y
<i>E. perriniana</i>	Spinning gum	Southeast Australia	...	Y	...	Y
<i>E. polyanthemus</i>	Red box	6	...	Southeast Australia	Y	Y
<i>E. torquata</i>	Coral gum	Western Australia	Y	Y
Nyssaceae s.s. (Cornaceae s.l.)								
<i>Camptotheca acuminata</i>	Happy tree	China and Tibet	Y	Y
Oleaceae								
<i>Chionanthus retusus</i>	Chinese fringetree	East Asia	Y	Y
<i>Fraxinus</i> sp.	Ash	Y
<i>F. uhdei</i>	Shamel ash	48	...	Central America	Y	Y	...	Y
<i>F. velutina</i>	Velvet ash	57	...	Southwestern North America	...	Y
<i>Olea europaea</i>	Olive	29	Y	Mediterranean, Asia and Africa	Y	Y	...	Y
<i>Osmanthus fragrans</i>	Sweet osmanthus	Himalayas to Japan	...	Y	Y	...
Onagraceae								
<i>Hauya microcerata</i>	Central America
Pinaceae								
<i>Cedrus atlantica</i>	Atlas cedar	1	...	Algeria and Morocco	...	Y	...	Y
<i>Keteleeria delavayi</i> (syn. <i>K. evelyniana</i>)	South China, Laos and Vietnam	...	Y
<i>Pinus densiflora</i>	Japanese red pine	Japan, Korea, China	...	Y	...	Y
<i>P. douglasiana</i>	Western Mexico	Y
Pittosporaceae								
<i>Hymenosporum flavum</i>	Native frangipani	20	...	Australia and New Guinea	Y	Y
<i>Pittosporum undulatum</i>	Victorian box	Eastern Australia	Y	Y	...	Y

(continued on next page)

HCO2198 (5'-TAAACTTCAGGGTGACCAAAAAATCA-3') (8). PCR and sequencing was performed as described by Rugman-Jones et al. (23). Sequences were aligned manually using BioEdit, version 7.0.9.0 (9). Primers were removed and each sequence was

translated (<http://www.ebi.ac.uk/Tools/emboss/transeq/index.html>) to confirm the absence of nuclear pseudogenes (26). All specimens were confirmed as PSHB (and not TSHB) by comparison with GenBank accession numbers JX912723–JX912725.

Table 1. (continued from preceding page)

Family, species ^b	Common name	N ^c	Crop ^d	Origin	FD ^e	S	P	G
Platanaceae								
<i>Platanus mexicana</i>	Mexican sycamore	Northeast Mexico	Y	Y
<i>P. occidentalis</i>	American sycamore	19	...	Eastern North America	Y	Y
* <i>P. racemosa</i>	California sycamore	68	...	California, Baja California	Y	Y
<i>P. wrightii</i>	Arizona sycamore	Southwest United States, Northwest Mexico	Y
Poaceae								
<i>Bambusa</i> sp.	Bamboo	Y
Podocarpaceae								
<i>Afrocarpus gracilior</i> (syn. <i>Podocarpus gracilior</i> , <i>Decussocarpus gracilior</i>)	Fern pine	East Africa
Proteaceae								
<i>Banksia saxicola</i>	Grampians banksia	Southeast Australia	Y	Y
<i>Macadamia integrifolia</i>	Macadamia nut	...	Y	Eastern Australia	Y	Y
Rhamnaceae								
<i>Rhamnus californica</i>	Coffee berry	Western North America	Y
<i>Ziziphus jujuba</i>	Chinese date	Temperate Asia	Y	Y
Rosaceae								
<i>Eriobotrya japonica</i>	Loquat	4	Y	China, Japan	Y
<i>Malus floribunda</i>	Japanese crabapple	2	Y	East Asia	...	Y	...	Y
<i>Prunus cerasoides</i>	Wild Himalayan cherry	South Asia	Y
<i>P. caroliniana</i>	Wild mock orange	Southeastern United States	...	Y	...	Y
<i>P. mume</i>	Chinese plum	Southern Japan	Y	Y
<i>P. persica</i>	Peach	4	Y	China	Y	Y	...	Y
<i>P. serrulata</i>	Japanese cherry	Japan, Korea and China	Y	Y
<i>Chaenomeles sinensis</i> (syn. <i>Cydonia sinensis</i>)	Chinese quince	China
<i>Pyrus kawakamii</i>	Evergreen pear	156	...	China, Taiwan	Y	Y
Rutaceae								
<i>Citrus sinensis</i>	Orange	6	Y	Southeast Asia	Y	Y	...	Y
Salicaceae								
<i>Salix babylonica</i>	Babylon willow	1	...	Asia	...	Y	Y	Y
<i>Salix</i> sp.	Y	Y	Y	Y
<i>Xylosma congestum</i>	Dense logwood	China	Y	Y
Sapindaceae								
<i>Alectryon excelsus</i>	Titoki	New Zealand	Y	Y	...	Y
<i>Harpullia arborea</i>	Tulip-wood tree	Tropical Southeast Asia	Y	Y	Y	
<i>H. pendula</i>	Tulipwood	Northeast Australia, Tropical Southeast Asia	...	Y	Y	Y
<i>Koelreuteria bipinnata</i>	Chinese flame tree	30	...	Southwest China	...	Y	...	Y
<i>K. elegans</i>	Chinese rain tree	Taiwan and Southern China	...	Y	...	Y
<i>K. elegans</i> subsp. <i>formsana</i> (syn. <i>K. henryi</i>)	Taiwan	...	Y	...	Y
<i>Ungnadia speciosa</i>	Mexican buckeye	South United States, North Mexico	Y	Y
Simaroubaceae								
<i>Ailanthus altissima</i>	Tree of heaven	2	...	China and Taiwan	Y
Taxodiaceae s.s. (Cupressaceae s.l.)								
<i>Metasequoia glyptostroboides</i> **	Dawn redwood	Western China	Y	Y
Theaceae								
<i>Camellia japonica</i>	Japanese camellia	Japan, Korea and China	Y	...
<i>Camellia</i> 'Apple Blossom'	Apple Blossom camellia	Horticultural origin	...	Y
<i>Camellia</i> 'Pink Sparkle'	Pink Sparkle camellia	Horticultural origin	...	Y
<i>C. reticulata</i>	Camellia	Southwest China	Y	Y
* <i>C. semiserrata</i>	Camellia	Y
<i>Cleyera japonica</i>	Sakaki	East Asia	Y	Y
Tiliaceae s.s. (Malvaceae s.l.)								
<i>Heliocarpus donnellsmithii</i>	Mexico and Central America	Y	Y
<i>Luehea divaricata</i>	Açoita-cavalo	Brazil	Y	Y	Y	Y
<i>Tilia americana</i>	Basswood	1	...	Central and Eastern North America	Y	Y	Y	...
Ulmaceae								
<i>Ulmus alata</i>	Winged elm	Central and Eastern United States	...	Y
<i>U. americana</i>	American elm	1	...	Eastern North America	Y
<i>U. parvifolia</i>	Chinese elm	183	...	East Asia	Y	Y
<i>Zelkova serrata</i>	Ju shu	1	...	East Asia	Y
Urticaceae								
<i>Pipturus argenteus</i>	Australia, Tropical Asia	Y	Y	Y	...
Verbenaceae								
<i>Aloysia virgate</i>	Sweet almond bush	South America
Vitaceae								
<i>Vitis vinifera</i>	Grapevine	...	Y	Mediterranean, Europe	Y	Y

Table 2. Apparent nonhosts of polyphagous shot hole borer

Family, species	Common name	Origin	N planted ^a
Aceraceae s.s. (Sapindaceae s.l.)			
<i>Acer rubrum</i>	Red maple	Eastern North America	1
Adoxaceae			
<i>Sambucus mexicana</i>	Mexican elderberry	Western North America	...
<i>Sambucus nigra</i>	Black elderberry	Europe, North Africa, Southwest Asia	...
Agavaceae s.s. (Asparagaceae s.l.)			
<i>Beaucarnea recurvata</i> (syn. <i>Nolina recurvata</i>)	...	Mexico	...
Aloeaceae s.s. (Xanthorrhoeaceae s.l.)			
<i>Aloe bainesii</i>	...	Southern Africa	...
<i>Aloe dichotoma</i>	...	Southern Africa	...
Anacardiaceae			
<i>Rhus lancea</i>	African sumac	South Africa	18
Apocynaceae			
<i>Acokanthera oppositifolia</i>	...	Tropical and Southern Africa	...
<i>Nerium oleander</i>	Oleander	Mediterranean, Middle and Central Asia	68
Areaceae			
<i>Archontophoenix cunninghamiana</i>	King palm	Eastern Australia	15
<i>Syagrus romanzoffianum</i> (syn. <i>Arecastrum romanzoffianum</i>)	Queen palm	South America	426
<i>Chamaerops humilis</i>	Mediterranean fan palm	Mediterranean	2
<i>Phoenix canariensis</i>	Canary Island date palm	Canary Islands	34
<i>P. dactylifera</i>	Date palm	Unknown origin	46
<i>P. reclinata</i>	Senegal date palm	Tropical Africa	...
<i>Trachycarpus fortunei</i>	Chinese windmill palm	Asia	80
Asteraceae			
<i>Montanoa guatemalensis</i>	...	Central America	...
Betulaceae			
<i>Betula nigra</i> 'Cully'	Heritage river birch	Horticultural origin	...
<i>Carpinus caroliniana</i>	America hornbeam	Eastern North America, Mexico, Central America	...
<i>Ostrya virginiana</i>	American hophornbeam	Eastern North America	...
Bignoniaceae			
<i>Jacaranda mimosifolia</i>	Jacaranda	South America	753
<i>Handroanthus chrysotrichus</i> (syn. <i>Tabebuia chrysotricha</i>)	Golden trumpet tree	Brazil	1
Boraginaceae			
<i>Ehretia latifolia</i> (syn. <i>E. austin-smithii</i>)	...	Mexico, Central America	...
Burseraceae			
<i>Bursera odorata</i>	...	Mexico	...
Cactaceae			
<i>Pereskia grandiflora</i>	...	Brazil	...
Caprifoliaceae			
<i>Weigela coraeensis</i>	Japanese weigela	Eastern Asia	...
Celastraceae			
<i>Euonymus bungeanus</i>	...	China, Korea	...
<i>Maytenus boaria</i>	Mayten	Chile	5
Cercidiphyllaceae			
<i>Cercidiphyllum japonicum</i>	Katsura tree	China, Japan	...
Clethraceae			
<i>Clethra macrophylla</i>	Mexican clethra	Mexico	...
Corynocarpaceae			
<i>Corynocarpus laevigatus</i>	Karaka	New Zealand	...
Cupressaceae			
<i>Calocedrus decurrens</i>	California incense cedar	Western North America	22
<i>Juniperus bermudiana</i>	Bermuda juniper	Bermuda	...
Ericaceae			
<i>Erica × darleyensis</i>	...	Horticultural origin	...
Fabaceae			
<i>Ceratonia siliqua</i>	Carob	Mediterranean, North Africa, Middle East and Western Asia	15
<i>Millettia reticulata</i>	Evergreen wisteria	South China, Taiwan	...
<i>Pithecellobium dulce</i>	Huamuchil	Mexico, Central America	...
<i>Sophora secundiflora</i>	Mountain laurel	Southwest United States, Mexico	...
Fagaceae			
<i>Lithocarpus glaber</i>	...	Japan, East China	...
<i>Quercus emoryi</i>	Emory oak	Southwest United States, Mexico	...
<i>Q. variabilis</i>	Chinese cork oak	China, Japan, Korea	...
Fouquieriaceae			
<i>Fouquieria macdougalii</i>	Mexican ocotillo	Northwest Mexico	...
Garryaceae			
<i>Garrya wrightii</i>	...	Southwest United States, Northwest Mexico	...
Ginkgoaceae			
<i>Ginkgo biloba</i>	Maidenhair tree	China	231

(continued on next page)

^a Number of times planted along streets of City of Orange (4).

Table 2. (continued from preceding page)

Family, species	Common name	Origin	N planted ^a
Hernandiaceae			
<i>Hernandia bivalvus</i>	Grease nut	Australia	...
Hippocastanaceae s.s. (Sapindaceae s.l.)			
<i>Aesculus hippocastanum</i>	Horse chestnut	Southeast Europe	2
<i>Aesculus × carnea</i>	Red horse chestnut	Horticultural origin	...
Iteaceae			
<i>Itea yunnanensis</i>	...	Southwest China	...
Lamiaceae			
<i>Vitex agnus-castus</i>	Vitex	Southern Europe	...
Lauraceae			
<i>Cryptocarya rubra</i>	...	Chile	...
Lythraceae			
<i>Lagerstroemia indica</i>	Crape myrtle	Asia	1,979
<i>L. subcostata</i>		Asia	...
<i>Punica granatum</i>	Pomegranate	Eastern Mediterranean to Himalayas	1
Magnoliaceae			
<i>Liriodendron tulipifera</i>	Tulip tree	Eastern North America	232
<i>Magnolia figo</i>	Banana shrub	China	...
Malvaceae			
<i>Brachychiton populneus</i>	Bottle tree tree	Northeast Australia	129
<i>Lagunaria patersonii</i>	Cow itch tree	Northeast Australia, South Pacific	1
Moraceae			
<i>Ficus carica</i>	Edible fig	Mediterranean to Near East	2
Myrtaceae			
<i>Acmena smithii</i>	Lilly pilli	Eastern Australia	1
<i>Agonis flexuosis</i>	Willow myrtle	Western Australia	...
<i>Angophora costata</i>	Smooth-bark apple	Eastern Australia	...
<i>Callistemon citrinus</i>	Lemon bottlebrush	Southeast Australia	433
<i>Eucalyptus cladocalyx</i>	Sugar gum	South Australia	8
<i>E. deglupta</i>	Rainbow gum	New Guinea to Philippines	...
<i>E. erythrocorys</i>	Red cap gum	Western Australia	5
<i>E. globulus</i>	Blue gum	Southeast Australia, Tasmania	43
<i>E. maculata</i>	Spotted gum	Eastern Australia	...
<i>E. microtheca</i>	Coolibah	Australia	...
<i>E. robusta</i>	Swamp mahogany	Northeast Australia	...
<i>E. saligna</i>	Sydney blue gum	Eastern Australia	...
<i>E. sideroxylon</i>	Red ironbark	Eastern Australia	9
<i>E. torrelliana</i>	Cadagah	Northeast Australia	...
<i>Acca sellowiana</i> (syn. <i>Feijoa sellowiana</i>)	Pineapple guava	Brazil, Argentina	...
<i>Lophostemon confertus</i>	Brush box	Eastern Australia	92
<i>Melaleuca lanceolata</i>	Moonah	South Australia	...
<i>M. linariifolia</i>	Snow-in-summer	Northeast Australia	...
<i>M. quinquenervia</i>	Cajeput	Eastern Australia	53
<i>Metrosideros excelsa</i>	New Zealand Christmas tree	New Zealand	...
<i>Syzygium buxifolium</i>	...	Chile	...
<i>S. paniculatum</i>	Brush cherry	Northeast Australia	...
Nyctaginaceae			
<i>Bougainvillea</i>	Bougainvillea
Nyssaceae s.s. (Cornaceae s.l.)			
<i>Nyssa sylvatica</i>	Sour gum	Eastern North America	...
Oleaceae			
<i>Ligustrum lucidum</i>	Chinese privet	East Asia	100
Papaveraceae			
<i>Bocconia arborea</i>	Tree poppy	Central America	...
Pinaceae			
<i>Pinus canariensis</i>	Canary Islands pine	Canary Islands	724
<i>P. halepensis</i>	Aleppo pine	Mediterranean	35
<i>P. pinea</i>	Italian stone pine	Mediterranean	11
<i>P. thunbergiana</i>	Japanese black pine	Japan, Korea	42
<i>P. torreyana</i>	Torrey pine	California	1
Pittosporaceae			
<i>Pittosporum tobira</i>	Japanese pittosporum	China, Japan	3
Poaceae			
<i>Phyllostachys aurea</i>	Golden bamboo	Southeast China	...
Podocarpaceae			
<i>Podocarpus elatus</i>	Rockingham podocarp	Eastern Australia	...
<i>P. macrophyllus</i>	Bigleaf podocarp	South China, Japan	2
<i>Nageia nagi</i> (syn. <i>Podocarpus nagi</i>)	Nagi	China, Taiwan, Japan	7
<i>P. totara</i>	Totara	New Zealand	...
Polygalaceae			
<i>Polygala apopetala</i>	Dwarf milkwort	Baja California	...
Proteaceae			
<i>Grevillea robusta</i>	Silky oak	Eastern Australia	9
<i>Stenocarpus sinuatus</i>	Firewheel tree	Eastern Australia	...

(continued on next page)

Table 2. (continued from preceding page)

Family, species	Common name	Origin	N planted ^a
Rhamnaceae			
<i>Hovenia dulcis</i>	Raisin tree	East Asia	...
<i>Ziziphus spina-christi</i>	...	East Africa, Middle and Near East	...
Rosaceae			
<i>Crataegus laevigata</i>	Midland hawthorn	Europe, North Africa, India	...
<i>Eriobotrya deflexa</i>	Bronze loquat	China, Taiwan	...
<i>Malus sylvestris</i>	Apple	Central Europe	1
<i>Photinia × fraseri</i>	Fraser's photinia	Horticultural origin	14
<i>Pyrus calleryana</i>	Ornamental pear	East Asia	2173
Rutaceae			
<i>Calodendrum capense</i>	Cape chestnut	South Africa	12
<i>Casimiroa edulis</i>	Cochitzapotl	Mexico and Central America	...
<i>Citrus limon</i>	Lemon	Unknown origin	5
<i>C. sinensis</i> 'Moro'	Moro blood orange	Horticultural origin	...
<i>Tetradium danieli</i> (syn. <i>Evodia danieli</i>)	...	China, Korea	...
<i>Geijera parviflora</i>	Australian willow	Australia	147
<i>Melicope elleryana</i>	Pink flowered soughwood	Australia	...
× <i>Citrofortunella</i>	...	Horticultural origin	...
Salicaceae			
<i>Dovyalis hebecarpa</i>	Ketembilla	Sri Lanka, Southern India	...
<i>Olmediella betschleriana</i>	Guatemalan holly	Mexico, Central America	...
<i>Oncoba spinosa</i>	Snuff-box tree	Africa, Arabia	...
Sciadopityaceae			
<i>Sciadopitys verticillata</i>	Japanese umbrella-pine	Japan	...
Scrophulariaceae			
<i>Eremophila bignoniflora</i>	Emu bush	Australia	...
<i>Myoporum laetum</i>	Myoporum	New Zealand	1
Solonaceae			
<i>Brugmansia</i> 'Charles Grimaldi'	Charles Grimaldi angel's trumpet	Horticultural origin	...
<i>Lycianthes rantonnei</i>	Blue potato bush	Argentina, Paraguay	1
Taxodiaceae s.s. (Cupressaceae s.l.)			
<i>Sequoia sempervirens</i>	Coast redwood	Western North America	...
<i>Taxodium mucronatum</i>	Mexican swamp cypress	Mexico	...
Theaceae			
<i>Camellia hiemalis</i> 'Kanjiro'	Kanjiro camellia	Horticultural origin	...
<i>C. sinensis</i>	Tea plant	Western China	...
<i>Gordonia axillaris</i>	...	Southeast Asia	...
Ulmaceae			
<i>Ulmus parvifolia</i>	Chinese elm	East Asia	175
Verbenaceae			
<i>Lippia umbellata</i> (syn. <i>L. torresii</i>)	...	Mexico, Central America	...

Table 3. Polyphagous shot hole borer (PSHB)–*Fusarium* sp. complex collected from tree species found in two botanical gardens located in a heavily infested part of Los Angeles County

Symptoms	Number of tree species	Fraction of all species (%)	Fraction of attacked species (%)
Total number	335
Attacked by PSHB	207	62	...
Host for <i>Fusarium</i> sp.	112	33	54
Weeping spots on bark	147	44	71
Powder depositions	22	7	11
Gumming on bark	69	20	33
Reproductive hosts for the beetle	19	6	9

Results

Host identification. Of 335 tree species (in 83 families) examined (Tables 1 and 2), 207 (62%) had symptoms consistent with attack by PSHB (small entry holes in tree bark, wet spots, gumming, white powdery exudate, and frass; Tables 1 and 3; Figs. 3A–K and 4A–C). The 207 species belonged to 58 families, with origins representing all the continents except Antarctica. Among the attacked species were 11 species native to California, and 13 that are agriculturally important crops in California (see below). There were 128 tree species in 58 families without obvious signs of attack by PSHB (Table 2). Thirty-three plant families examined contained species that were both attacked and not attacked by PSHB.

Fungal DNA was extracted from *Fusarium* sp. isolates recovered from symptomatic tissues of 113 of the 207 tree species (54%) attacked by PSHB. In all cases, sequences of the ITS, EF1- α , RPB1, and RPB2 genes of the fungus were identical to the respec-

tive known sequences of isolates of *Fusarium* sp. from Mendel et al. (15) (GenBank accessions JQ038007 to -038034) and Eskalen et al. (7) (GenBank accessions JQ723753 to -723756 and JQ723760 to -723763). These findings indicate the susceptibility of the host tree to the growth of *Fusarium* sp., at least at the site of the entry hole. Fungal sequences generated in the current study have been deposited in GenBank under the following accession numbers: ITS, JX891673 to -JX891784; EF1- α , JX891785 to -JX891896; RPB1, JX891897 to -JX892008; and RPB2, JX892009 to -JX892120. *Fusarium* sp. was isolated from 113 tree species in 40 families. Families with the highest number of infected species were the Aceraceae, Fabaceae, and Fagaceae.

Of the tree species susceptible to varying degrees of attack by PSHB, 19 were suitable for beetle reproduction. Suitability for reproduction was confirmed by either the presence of eggs, larvae, pupae, or teneral females or the presence of males in the galleries (Table 3; Fig. 5). This number might be an underestimate, because




Fig. 3. A–D, External symptoms of staining; E–H, gumming; and I–K, branch dieback for polyphagous shot hole borer (PSHB)–*Fusarium* dieback disease on **A**, *Quercus agrifolia*; **B**, *Liquidambar styraciflua*; **C**, *Platanus racemosa*; **D**, *Senna specatibilis* var. *excelsa* (syn. *Cassia excelsa*); **E**, *Alectryon excelsus*; **F**, *Acacia melanoxylon*; **G**, *Alianthus altissima*; **H**, *Koelreuteria elegans*; and **I–K**, *Persea americana*.

confirmation requires cutting into trees, which was not logistically possible at the time due to the limited number of trees that represent each species within the botanical gardens' collection.

All tree species considered to be reproductive hosts had severe branch dieback (Fig. 3I–K), and the death of mature reproductive hosts that were infested with FD was observed for *Quercus robur*, *Acer negundo*, and *R. communis*. *A. negundo* and *Q. robur* also exhibited leaf wilting and discoloration on the branches prior to dieback and tree death.

Symptom characterization. Symptoms surrounding the beetle entry hole included staining (71% of hosts), gum deposition (33% of hosts), and white powder-like exudates (11% of hosts) (Figs. 3, 4, and 6, Tables 1 and 3). Staining was characterized by a moist discoloration of the outer bark that oozed out of the entry holes in various amounts, depending on the host (Fig. 3A–D). Discoloration caused by fungal colonization was observed underneath the

inner bark layer when the beetle entry hole penetrated through the cambium and into the xylem (Figs. 7 and 8). On various hosts, PSHB attack resulted in the deposition of an amber colored gum that exuded through the entry holes and varied in consistency between spongy and pliable to solid and hard, depending on the host (Table 1, Fig. 3E–H). After peeling back the outer bark beneath the gum deposition, tissue discoloration was observed and subsequent recovery of *Fusarium* sp. occurred on 46% of the tree species showing a gumming response (Table 2). Otherwise, the beetle was found embedded within the gum deposit, and there was no discoloration of the plant tissue. A white powdery exudate was observed on some species (Table 1). On avocado, it characteristically emerged from entry holes and formed a white “sugar volcano” approximately 1 to 15 cm in diameter and 0.5 to 5 cm in height (Fig. 4A). Frass was occasionally observed, particularly on trees in the Aceraceae (Fig. 4B).




Fig. 4. A, External symptoms of polyphagous shot hole borer and *Fusarium* dieback disease that are characteristic for *Persea americana* (white powder exudate/sugar volcano); B, *Acer negundo*, (abundant entry holes associated with frass); and C, *Ricinus communis* (staining).




Fig. 5. Offspring (top arrow) and larvae (bottom arrow) in beetle galleries confirm *Acer negundo* as a reproductive host of polyphagous shot hole borer.

In total, 26% of the identified hosts of FD had branch dieback on over 50% of the tree. Symptoms of branch dieback were typically observed on primary branches where areas of weakness caused by beetle and fungal colonization were observed at the branch nodes (Figs. 3I–K and 8A–F). For trees attacked by PSHB, the likelihood of a higher than 50% branch dieback was positively correlated with presence of *Fusarium* sp. and increasing severity of PSHB attack, and negatively correlated with symptoms of gumming (Table 4).

Estimated frequency of trees affected by PSHB–*Fusarium* sp. complex in the urban forest. The botanical garden survey was used to determine the susceptibility of 103 common street tree species (Table 5). In all, 53 species were attacked by PSHB and, of these, *Fusarium* sp. was isolated from 36. The beetle was able to reproduce in six of these species. Relating these numbers to the relative abundance of the different trees in the urban environment of southern California (4) reveals that, of the individual street trees, over half (56%) belong to species that are attacked by the PSHB,




Fig. 6. Proportion of trees prone to polyphagous shot hole borer attack with respective symptoms.

48% belong to species that are hosts to the *Fusarium* sp., and 26% belong to species that are reproductive hosts of PSHB (Table 5).

FD complex in agriculturally important plants. In this survey, 13 agriculturally important crops displayed signs and symptoms consistent with PSHB attack (Table 1). Of these, growth of the *Fusarium* sp. was confirmed on avocado, eastern mulberry, olive, macadamia, Turkish hazelnut, loquat, peach, grapevine, and citrus (although the latter does not appear to be commonly attacked). Avocado was the only species confirmed as a reproductive host for PSHB. Symptoms consistent with attack by PSHB were detected on cassava, Japanese persimmon, Japanese crabapple, and eastern black walnut but the *Fusarium* sp. was not detected growing on these species. The following agricultural crops showed no evidence of PSHB attack: date palm, pomegranate, edible fig, apple, and most citrus tree varieties (see above; Table 2).

Discussion

Host range, symptoms, and plant responses. In this study, we present the host range of an emerging generalist pest–pathogen complex in California. The behavior of PSHB was consistent with



Fig. 7. Evidence of an unsuccessful attempt by polyphagous shot hole borer to establish a gallery in *Acer pectinatum* subsp. *laxiflorum*. A, Depth of penetration by the beetle is less than 1 mm. B, No sign of fungal colonization beneath the cambium.




Fig. 8. A, Internal symptoms of polyphagous shot hole borer or *Fusarium* dieback disease showing gallery formation discoloration caused by fungal colonization on *Quercus agrifolia*; B, *Persea americana*; C, *Quercus robur*, and D, *Ricinus communis* in cross section; and E, *Acer negundo* and F, *R. communis* in longitudinal section.

the ecological group of first-degree primary polyphagous borers that invade healthy trees in many different plant families (5,12,22,24). Plant species attacked by the FD disease complex have worldwide origins, and include species such as the critically endangered dawn redwood (*Metasequoia glyptostroboides*) from China. In purely economic terms, however, the threat posed by FD to important agricultural crops, ornamental trees used for landscaping, and native forest tree species in California is a major concern. The wide host range of PSHB described in our southern California study was consistent with literature reports for the “tea shot hole borer” (5). However, with the present uncertainty about the species status of individuals morphologically recognized as *E. formicatus*, the host range reported by Danthanarayana (5) may actually refer to the hosts of two (or more) cryptic species, each with a more limited host range.

Currently, it is difficult to estimate the longevity of trees infested by either the beetle or fungus alone, or trees that are reproductive

hosts. Even though a tree species is not a reproductive host, an attempted PSHB attack may serve as an infection site for *Fusarium* sp. In some cases, *Fusarium* sp. was not able to infect the tissue. Even on known hosts of the fungus, tree infection required penetration into at least the cambium layer, which generally varied in depth below the outer bark between species. Given that 82% of species susceptible to FD in the present study are not reproductive hosts, we conclude that infection is most likely due to susceptibility of the tree to the fungus if the beetle is able to penetrate into or through this critical layer of tissue. Based on the previous observations of *Fusarium* sp. clogging conductive tissue within the tree (7,15) and dead or dying reproductive hosts infested with FD, and given that there is a correlation between severity of the beetle attack (which thereby increases severity of infection by *Fusarium* sp.) and the observed dieback, we may predict that, over time, infection by *Fusarium* sp. may lead to tree death in some species.

Although gumming was present on some tree species from which *Fusarium* sp. was also recovered, severe branch dieback was more likely in trees that did not exhibit gumming. This suggests that trees with gumming may be more resistant to beetle (12) and fungal attack. Most trees with gumming did not have necrotic tissues beneath the bark because the gum provided a physical barrier for beetle and fungal entry into the plant. This pattern is consistent with attack to trees by the TSHB in southern Asia, although there is no mention of resistance to fungal infection (5). Host response and resistance to disease or beetle infestation of this particular complex needs to be studied further to control disease establishment and spread.

Potential threat to California landscapes, agriculture, and global plant communities. The research and understanding of the distribution and impact of the FD complex are in the very early stages. However, the ability of this complex to attack multiple tree species in a large number of families suggests that it may become established across a range of plant communities on both a local and global scale. *Phytophthora ramorum* is an example of a generalist pathogen within California that also has a global distribution. It is the cause of sudden oak death, or ramorum dieback. This pathogen has killed thousands of oaks, tanoaks, and other woody hosts in California plant communities (6,13,19,21). However, the number of families susceptible to natural attack by *Fusarium* sp. in this study is five times higher than that of *P. ramorum* (6).

Reproductive hosts are negatively impacted when infected with FD and perpetuate healthy populations of PSHB, which facilitates long-term disease establishment and spread within a system. This study revealed that 5 of the 11 California native plant species surveyed were reproductive hosts of PSHB. This suggests that native plant communities and landscapes with these host plants are in danger of attack. Natural plant communities that are particularly vulnerable in southern California because reproductive hosts of PSHB are dominant members of the plant community include mixed evergreen forests, oak woodlands, foothill woodlands, and riparian habitats. Also noteworthy was that each continent on the globe had at least one reproductive host in this study, excluding Antarctica. This indicates a potential for FD establishment on a global scale. Studies on climatic conditions and other factors that influence PSHB distribution need to be implemented to determine the degree to which areas in other parts of the world are at risk for establishment by PSHB and *Fusarium* sp.

The juxtaposition of agriculture, urban landscapes, and forests in southern California harboring reproductive host species allows for potential rapid disease spread over a large geographic area that would otherwise be limited to islands of suitable habitat for PSHB and *Fusarium* sp. As an example, the urban forest in southern Cali-

Table 4. Parameter estimate, odds ratio, and *P* values for the logistic regression model describing factors contributing to the likelihood of branch dieback >50% on trees attacked by the polyphagous shot hole borer (PSHB) (whole model $P > \chi^2$ 0.0035 and 0.0003)

Variable	Estimate	Odds ratio	$P > \chi^2$
<i>Fusarium</i> sp.	0.5269	2.869	0.0065
Gumming	-0.4048	2.610	0.0202
PSHB severity	0.9553	3.4147–6.9055	0.0004

Table 5. Number of tree species and abundance of tree specimens susceptible to the polyphagous shot hole borer (PSHB)–*Fusarium* sp. complex in the southern Californian urban forest^a

Variable	Number of species	Abundance (%) ^b
Total number	103	...
Attacked by PSHB	53	56
Host for <i>Fusarium</i> sp.	36	48
Reproductive hosts for the beetle	6	26

^a Estimations were made using a representative tree species list (4).

^b Percentage of all specimens in the urban forest.

fornia could exacerbate the spread of FD, given that 48% of the street trees in southern California (4) belong to species that were confirmed as susceptible to FD. Six of these tree species are also reproductive hosts for PSHB, including American sweetgum, which is one of the most commonly planted street trees. Additionally, castor bean, a reproductive host and weed with a global distribution, is prevalent in urban, agriculturally important, and riparian areas and in the interface between these landscapes in southern California (27). The potential costs associated with the infestation of the urban forest may become substantial.

The FD complex is of special concern to the California avocado industry. It is already established and causing significant losses in commercial avocado groves in Israel (15). Given the high frequency and broad distribution of reproductive hosts and other host species throughout the region, disease spread and widespread establishment into commercial avocado groves in southern California is almost inevitable. At the time of writing, symptoms of FD have not been observed in commercial citrus groves or vineyards but have been confirmed in two commercial avocado groves in Los Angeles County. Disease distribution throughout southern California is currently being investigated.

Although studies are underway to determine host resistance, disease distribution, and impact, the results from this study suggest that plants confirmed as nonhosts, or hosts that display symptoms of resistance to PSHB, be given priority in future landscape plantings.

Acknowledgments

We thank the California Avocado Commission for supporting this research; Huntington Library, Art Collections, and Botanical Gardens and the Los Angeles County Arboretum for allowing us to conduct the survey for this study; P. Meyers from the Huntington herbarium for assistance with editing the scientific names of host and nonhost species; F. McDonogh for assistance with surveying the LA Arboretum; J. Mayorquin, D. Wang, L. Yeung, and F. Na (Department of Plant Pathology and Microbiology, University of California, Riverside) for assistance with sample processing and molecular identification of fungi; A. Coelho, Jr. (Department of Entomology, University of California, Riverside) for assistance with sample collection; R. Hofshi and M. L. Arpaia for ongoing support in this research; and the anonymous reviewers of this manuscript for comments and suggestions.

Literature Cited

- Benson, D. A., Karsch-Mizrachi, I., Lipman D. J., Ostell J., and Wheeler D. L. 2008. GenBank. *Nucleic Acids Res* 36:D25–D30.
- Cenis, J. L. 1992. Rapid extraction of fungal DNA for PCR amplification. *Nucleic Acids Res.* 20:2380.
- Chang, D. C., G. B. Grant, K. O'Donnell, K. A. Wannemuehler, J. Noble-Wang, C. Y. Rao, L. M. Jacobson, C. S. Crowell, R. S. Sneed, F. M. T. Lewis, J. K. Schaffzin, M. A. Kainer, C. A. Genese, E. C. Alfonso, D. B. Jones, A. Srinivasan, S. K. Fridkin, and B. J. Park. 2006. A multistate outbreak of *Fusarium* keratitis associated with use of a new contact lens solution. *JAMA* 296:953–963.
- City of Orange Species Frequency Report with Characteristics website. 1999. <http://www.cityoforange.org/civifax/filebank/blobdload.aspx?blobid=3499>
- Danthanarayana, W. 1968. The distribution and host-range of the shot-hole borer (*Xyleborus formicatus* Eichh.) of tea. *Tea Q.* 39:61–69.
- Davidson, J. M., Werres, S., Garbelotto, M., Hansen, E. M., and Rizzo, D. M. 2003. Sudden oak death and associated diseases caused by *Phytophthora ramorum*. *Plant Health Prog.* Online publication. doi:10.1094/PHP-2003-0707-01-DG
- Eskalen, A., Gonzalez, A., Wang, D. H., Twizeyimana, M., Mayorquin, M., and Lynch, S. C. 2012. First report of *Fusarium* sp. and its vector tea shot hole borer (*Euwallacea formicatus*) causing Fusarium dieback on avocado in California. *Plant Dis.* 96:1070.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., and Vrijenhoek, R. 1994. DNA primers for amplification of mitochondrial cytochrome C oxidase subunit I from diverse metazoan invertebrates. *Mol. Marine Biol. Biotechnol.* 3:294–299.
- Hall, T. A. 1999. BioEdit: a user friendly biological sequence alignment and analysis program for Windows 95/98/NT. *Nucleic Acids Symp.* 41:95–98.
- Hebert, P. D. N., Cywinska, A., Ball, S. L., and deWard, J. R. 2003. Biological identifications through DNA barcodes. *Proc. R. Soc. Lond. B* 270:313–321.
- Hulcr, J., Mann, R., and Lukasz, L. S. 2011. The scent of a partner: ambrosia beetles are attracted to volatiles from their fungal symbionts. *J. Chem. Ecol.* 37:1374–1377.
- Kovach, J., and Gorsuch, C. S. 1985. Survey of ambrosia beetle species'

- infesting South Carolina peach orchards and a taxonomic key for the most common species. *Agric. Entomol.* 2:238-247.
13. Maloney, P. E., Lynch, S. C., Kane, S. F., Jensen, C. E., and Rizzo, D. M. 2005. Establishment of an emerging generalist pathogen in redwood forest communities. *J. Ecol.* 93:899-905.
 14. McDonald, V., and Eskalen, A. 2011. Botryosphaeriaceae species associated with avocado branch cankers in California. *Plant Dis.* 95:1465-1473.
 15. Mendel, Z., Protosav, A., Sharon, M., Zveibil, A., Ben Yahuda, S., O'Donnell, K., Rabaglia, R., Wysoki, M., and Freeman, S. 2012. An Asian ambrosia beetle *Euwallacea fornicatus* and its novel symbiotic fungus *Fusarium* sp. pose a serious threat to Israeli avocado industry. *Phytoparasitica*, Online publication. doi:10.1007/s12600-012-0223-7
 16. O'Donnell, K., Sarver, B. A. J., Brandt, M., Chang, D. C., Noble-Wang, J., Park, B. J., Sutton, D. A., Benjamin, L., Lindsley, M., Padhye, A., Geiser, D. M., and Ward, T. J. 2007. Phylogenetic diversity and microsphere array-based genotyping of human pathogenic fusaria, including isolates from the multistate contact lens-associated U.S. keratitis outbreaks of 2005 and 2006. *J. Clin. Microbiol.* 45:2235-2248.
 17. O'Donnell, K., Cigelnik, E., and Nirenberg, H. 1998. Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* 90:465-493.
 18. O'Donnell, K., Sutton, D. A., Rinaldi, M. G., Sarver, B. A. J., Arunmozhi Balajee, S., Schroer, H., Summerbell, R. C., Robert, V. A. R. G., Crous, P. W., Zhang, N., Aoki, T., Jung, K., Park, J., Lee, Y., Kang, S., Park, B., and Geiser, D. M. 2010. Internet-accessible DNA sequence database for identifying *Fusarium* from human and animal infections. *J. Clin. Microbiol.* 48:3708-3718.
 19. Parke, J. L., and Lucas, S. 2008. Sudden oak death and ramorum blight. *Plant Health Instructor*. Online publication. doi:10.1094/PHI-I-2008-0227-01
 20. Rabaglia, R. J., Dole, S. A., and Cognato, A., 2006. Review of American *Xyleborina* (Coleoptera: Curculioniadae: Scolytinae) occurring north of Mexico, with an illustrated key. *Ann. Entomol. Soc. Am.* 99:1034-1055.
 21. Rizzo, D. M., and Garbelotto, M. 2003. Sudden oak death: endangering California and Oregon forest ecosystems. *Front. Ecol. Environ.* 1:197-204.
 22. Rudinsky, J. A. 1962. Ecology of Scolytidae. *A. Rev. Ent.* 7:327-348.
 23. Rugman-Jones, P. S., Hoddle, M. S., Amrich, R., Heraty, J. M., Stouthamer-Ingel, C. E., and Stouthamer, R. 2012. Phylogeographic structure, outbreeding depression, and reluctant virgin oviposition in the bean trips, *Caliothrips fasciatus* (Pergande) (Thysanoptera: Thripidae), in California. *Bull. Entomol. Res.* Online publication. doi:10.1017/S0007485312000302
 24. Schedl, K. E. 1958. Breeding habits of Arboricole insects in central Africa. Pages 183-197 in: *Proc. Int. Cong. Ent.* 10th Meet. Montreal, Quebec, Canada.
 25. Slippers, B., Crous, P. W., Denman, S., Coutinho, T. A., Wingfield, B. D., and Wingfield, M. J. 2004. Combined multiple gene genealogies and phenotypic characters differentiate several species previously identified as *Botryosphaeria dothidea*. *Mycologia* 96:83-101.
 26. Song, H., Buhay, J. E., Whiting, M. F., and Crandall, K. A. 2008. Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proc. Natl. Acad. Sci. USA* 105:13486-13491.
 27. United States Department of Agriculture, Natural Resources Conservation Service. 2012. Plant Profile website. <http://plants.usda.gov/java/profile?symbol=RICO3>
 28. Wood, S. L. 1989. Nomenclatural changes and new species of Scolytidae (Coleoptera), Part IV. *Great Basin Nat.* 49:167-185.
 29. Zhang, N., O'Donnell, K., Sutton, D. A., Nalim, F. A., Summerbell, R. C., Padhye, A. A., and Geiser, D. M. 2006. Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. *J. Clin. Microbiol.* 44:2186-2190.